



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ger, of Nashville, Tenn. A few of them may here be put on record. *Forestiera ligustrina* does not (like *F. acuminata*) blossom in early spring from axils of the preceding year. On the cedar barrens near Nashville, where it abounds, the fragrant flowers develop about the middle of August from the axils of the leaves of that year; and the fruit ripens at the end of September. That of *F. acuminata*, which blossoms very early, is ripe before the end of May. *Tragia macrocarpa*.—Both surfaces of the leaves bear stinging hairs. *Phlox Stellaria*.—This neat and rare species is found at Lavergne, seventeen miles from Nashville, in cedar barrens, growing in beds of sphagnum and other mosses, in moist places. *Thermopsis Caroliniana*, a most rare species, has been found by Dr. Gottinger on the Harpeth hills, near Nashville.—A. GRAY.

LITTORELLA AND SCHIZÆA IN NOVA SCOTIA.—It is singular how a long-overlooked plant, once detected, is then promptly found again and again. Following upon Mr Pringle's announcement of the second discovery of *Littorella lacustris* (at the northern end of Lake Champlain), I have now to announce that Miss Elizabeth G. Knight, of the New York Normal College, found it in August last, growing abundantly between the stones on the shores of Grand Lake, Nova Scotia, twenty-three miles from Halifax.

Botanists will be equally interested to know that she also detected, among the rhizomes of *Osmunda regalis*, near the lake shore, the rare *Schizæa pusilla*. La Pylaie's specimens in his herbarium at Paris, collected in Newfoundland about sixty years ago (which I have seen), had accredited this plant to New Foundland; but I believe no one has since found it out of New Jersey until this happy discovery by Miss Knight in an adjacent portion of British America. A. GRAY.

THE COEFFICIENT OF CONTRACTION.—My attention was lately drawn to a remarkable case of the difference in length produced by unequal "seasoning," in the two sides of an oak post. The post referred to is about four inches square; one end is sunk in the ground and the other projects ten and one half feet above the surface. When placed in position some time ago it was straight and perpendicular; at the present time it leans toward the south, deviating a little over a foot from the perpendicular. The post was "set with the compass," and it is interesting to note that a north and south line lies in the plane of the curve.

Experiments have given us the coefficients of expansion in different metals. Why may we not have experiments to determine the coefficients of contraction in different woods, i. e., to determine the fractional decrease in length produced in rods of "green" wood, say 1 meter long and 2 cm. square, by the application of a steady and absolute dry heat for a given length of time? The knowledge would be of no practical importance perhaps, but it might bring out unsuspected correlation between looseness of tissue and amount of contraction.—C. R. BARNES, *La Fayette, Ind.*